

Behavioral Risk Factor Surveillance System (BRFSS) Health Indicators by County

Data from the 2016 Behavioral Risk Factor Surveillance System were used to generate percentages of non-institutionalized adult (18+) NYS residents for various health indicators for a range of geographies.

<https://health.data.ny.gov/Health/Behavioral-Risk-Factor-Surveillance-System-BRFSS-H/isy7-eb4n>

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury	Franklin							rates suppressed due to small sample size
2016	Percentage of adults reporting a depressive disorder	Franklin	16.3	11.7	20.8	17.6	12.3	22.8	
2016	Percentage of adult binge drinking during the past month	Franklin	17.7	12.3	23	17.8	12.4	23.3	
2016	Percentage of adults aged 18+ years with flu immunization in the past year	Franklin	31.2	25.5	36.9	29.3	24	34.6	
2016	Percentage of adults aged 18-24 years who are current smokers	Franklin							rates suppressed due to small sample size
2016	Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year	Franklin	63.8	55.5	72.2				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 18-64 years with healthcare coverage	Franklin	88.3	81.6	95				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 50-64 years receiving colorectal cancer screening based on recent guidelines	Franklin	66.3	57.1	75.5				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	Franklin	74.1	67.4	80.7				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with at least one fall in the past 12 months	Franklin	26.7	18.7	34.6				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with flu immunization in the past year	Franklin	59.1	49.7	68.6				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with pneumococcal immunization	Franklin	79.1	71.6	86.6				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults experiencing housing insecurity in the past 12 months	Franklin	22.9	16.3	29.5	23.7	15.1	32.3	
2016	Percentage of adults overweight or obese	Franklin	71.3	64.7	77.9	72.2	65.5	78.9	
2016	Percentage of adults reporting 14 or more days of poor physical health	Franklin	13.5	8.9	18.1	12.1	8.3	15.9	
2016	Percentage of adults reporting driving under the influence in the past 30 days	Franklin							rates suppressed due to small sample size
2016	Percentage of adults reporting poor self-reported health	Franklin	5.2	1.7	8.8	4.6	1.7	7.4	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults who are current smokers	Franklin	28.8	22.1	35.6	29.5	22.3	36.8	
2016	Percentage of adults who are obese	Franklin	35.9	29.4	42.3	34.8	27.8	41.9	
2016	Percentage of adults who consume one or more sugary drinks daily	Franklin	24.2	18.7	29.8	24.5	18.5	30.5	
2016	Percentage of adults who did not receive medical care because of cost	Franklin	7.5	3.5	11.6	7.1	3.4	10.7	
2016	Percentage of adults who had a dentist visit within the past year	Franklin	57.2	50.4	64	56	48.7	63.4	
2016	Percentage of adults who had a test for high blood sugar or diabetes within the past three years	Franklin	58.1	50.3	65.8	60.4	53.3	67.5	
2016	Percentage of adults who have a regular health care provider	Franklin	78.9	72.3	85.5	81.1	76.3	85.9	
2016	Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition	Franklin	7.6	4.1	11	4.9	2.7	7	
2016	Percentage of adults who participated in leisure time physical activity in the past 30 days	Franklin	72.1	66.2	78.1	72.1	65.5	78.8	
2016	Percentage of adults who report consuming less than one fruit and less than one vegetable daily (no fruits or vegetables)	Franklin	35.7	28.6	42.9	36.6	28.8	44.4	
2016	Percentage of adults who use e-cigarettes	Franklin	5.2	1.8	8.5	6.3	1.9	10.7	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Percentage of adults with an annual household income less than \$25,000 who are obese	Franklin	26.8	16.8	36.9	26	16.8	35.2	unreliable crude rate due to large standard error
2016	Percentage of adults with annual household income less than \$25,000 who are current smokers	Franklin	39.7	25.4	54.1	41.9	30.1	53.8	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with arthritis	Franklin	35.5	29.1	41.8	33.6	27.6	39.7	
2016	Percentage of adults with cardiovascular disease (heart attack, coronary heart disease, or stroke)	Franklin	11.4	7.4	15.4	9.8	6.6	12.9	
2016	Percentage of adults with COPD	Franklin	8.5	5.6	11.3	7.7	5.1	10.4	
2016	Percentage of adults with current asthma	Franklin	13.7	8.9	18.4	13.8	8.7	18.9	
2016	Percentage of adults with diagnosed high blood pressure taking high blood pressure medication	Franklin	74.3	63.7	85	50.2	37	63.4	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with disability (based on 6 ACA disability questions)	Franklin	24.5	18.9	30.2	21.6	16.6	26.5	
2016	Percentage of adults with disability who are obese	Franklin	41.6	29.2	54.1				unreliable crude rate due to large standard error; age-adjusted rate not defined due to empty age-adjustment cells
2016	Percentage of adults with disability who smoke	Franklin	46.9	33.6	60.1				unreliable crude rate due to large standard error; age-adjusted rate not defined due to empty age-adjustment cells
2016	Percentage of adults with disability with cardiovascular disease (heart attack, coronary heart disease, or stroke)	Franklin	20.3	10.9	29.6				age-adjusted rate not defined due to empty age-adjustment cells
2016	Percentage of adults with food security over the last 12 months	Franklin	81.6	75.2	88	81.5	75.4	87.7	
2016	Percentage of adults with physician diagnosed diabetes	Franklin	12.8	9.4	16.2	11.1	8.4	13.7	
2016	Percentage of adults with physician diagnosed high blood pressure.	Franklin	36.1	29.9	42.4	32	27	36.9	
2016	Percentage of adults with physician diagnosed prediabetes	Franklin	7.7	4.9	10.6	7.8	5	10.5	
2016	Percentage of adults with poor mental health for 14 or more days in the last month	Franklin	12.5	7.7	17.3	13.1	7.6	18.6	
2016	Percentage of cigarette smoking among adults who report poor mental health	Franklin	36.1	14.6	57.6	35.2	18	52.4	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of men aged 55-69 receiving shared decision making for PSA screening (discussed the advantages and disadvantages of the PSA test with a health professional)	Franklin	34.5	22	47				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-44 years who had dental visit within the last year	Franklin	57.9	42.1	73.7				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy	Franklin	44.9	28.2	61.6				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-64 years with health care coverage	Franklin	94.2	89	99.4				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 21-29 years receiving cervical cancer screening based on 2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test plus HPV test within past five years for women aged 30-65)	Franklin							rates suppressed due to small sample size
2016	Percentage of women aged 21-65 years receiving cervical cancer screening based on 2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test plus HPV test within past five years for women aged 30-65)	Franklin	80	68.2	91.8				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines	Franklin	78.9	70.6	87.2				age-adjusted rate not defined for indicators with an age restriction
2016	Women aged 18-44 years who saw a doctor for a routine checkup within the last year	Franklin	62.7	47.5	78				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction