## Behavioral Risk Factor Surveillance System (BRFSS) Health Indicators by County

Data from the 2016 Behavioral Risk Factor Surveillance System were used to generate percentages of non-institutionalized adult (18+) NYS residents for various health indicators for a range of geographies.

https://health.data.ny.gov/Health/Behavioral-Risk-Factor-Surveillance-System-BRFSS-H/jsy7-eb4n

			Crude	CL 95% Lower,	CL 95% Upper,		CL 95% Lower, Age	-CL 95% Upper, Age	
Year	Health Indicator	County	Rate	Crude	Crude	Age-adjusted Rate	adjusted	adjusted	Notes
									unreliable crude rate due to large standard error; age-
	Among adults aged 65+ years with at least one fall in the past 12 months, percent that								adjusted rate not defined for indicators with an age
2016	resulted in injury	St. Lawrence	33.7	19.7	47.8	В			restriction
2016	Pecentage of adults reporting a depressive disorder	St. Lawrence	15.9	11	20.8	16.9	11.5	22.2	
2016	Percentage of adult binge drinking during the past month	St. Lawrence	19.4	13.1	25.8	20	13.3	26.7	
2016	Percentage of adults aged 18+ years with flu immunization in the past year	St. Lawrence	38.3	31.7	7 44.9	36.3	29.4	43.1	
2016	Percentage of adults aged 18-24 years who are current smokers	St. Lawrence							rates suppressed due to small sample size
	Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within								age-adjusted rate not defined for indicators with an age
2016	the last year	St. Lawrence	71.5	63.5	79.4				restriction
	·								age-adjusted rate not defined for indicators with an age
2016	Percentage of adults aged 18-64 years with healthcare coverage	St. Lawrence	97.2	95	99.4				restriction
	Percentage of adults aged 50-64 years receiving colorectal cancer screening based on								age-adjusted rate not defined for indicators with an age
2016	recent guidelines	St. Lawrence	78.5	70.7	86.3	s			restriction
	Percentage of adults aged 50-75 years receiving colorectal cancer screening based on								age-adjusted rate not defined for indicators with an age
2016	recent guidelines	St. Lawrence	79.5	73.3	85.7	,			restriction
									age-adjusted rate not defined for indicators with an age
2016	Percentage of adults aged 65+ years with at least one fall in the past 12 months	St. Lawrence	38	28.8	47.3	3			restriction
	,								age-adjusted rate not defined for indicators with an age
2016	Percentage of adults aged 65+ years with flu immunization in the past year	St. Lawrence	66.9	57.7	76.1				restriction
2010	refeetinge of dudies aged ost years with his minimum and of the past year	St. Edwichee	00.5	37.7	70.3				age-adjusted rate not defined for indicators with an age
2016	Percentage of adults aged 65+ years with pneumococcal immunization	St. Lawrence	74.8	66.2	83.5				restriction
2010	Percentage of addits aged 05+ years with phedinococcar initiditization	3t. Lawrence	74.0	00.2	03.3	1			restriction
2016	Percentage of adults experiencing housing insecurity in the past 12 months	C+ Lauranca	30.2	23.3	37.1	31.4	23.3	39.6	
2016	Percentage of adults experiencing housing insecurity in the past 12 months  Percentage of adults overweight or obese	St. Lawrence St. Lawrence	68.5				62		I.
2016	Percentage of adults overweight or obese  Percentage of adults reporting 14 or more days of poor physical health	St. Lawrence	13.5				8.9		
2010	Percentage of addits reporting 14 or more days of poor physical health	St. Lawrence	15.3	9.2	17.0	12.0	0.9	10.2	
									unreliable crude rate due to large standard error;
2016	Percentage of adults reporting driving under the influence in the past 30 days	St. Lawrence	4.2	0.9	7.5		1.1	7.4	unreliable age-adjusted rate due to large standard error
2016	Percentage of addits reporting driving under the initiaerice in the past 50 days	St. Lawrence	4.2	0.5	7.5	4.3	1.1	7.4	unrenable age-adjusted rate due to large standard error
2016	Davagata as of adulta associate associate associated baselik	St. Lawrence	4.3	1.9		4.3	1.7	_	
	Percentage of adults reporting poor self-reported health		14.4				1.7		unreliable age-adjusted rate due to large standard error
2016	Percentage of adults who are current smokers	St. Lawrence					26.6		
2016 2016	Percentage of adults who are obese	St. Lawrence	34.2 27.5				26.6		
	Percentage of adults who consume one or more sugary drinks daily	St. Lawrence							
2016	Percentage of adults who did not receive medical care because of cost	St. Lawrence	6.3	55.6			2.6 55.3	69.5	unreliable crude rate due to large standard error
2016	Percentage of adults who had a dentist visit within the past year	St. Lawrence	62.4	55.6	69.2	62.4	55.3	69.5	
2046	Percentage of adults who had a test for high blood sugar or diabetes within the past	Ct. Lawrence			,				
2016	three years	St. Lawrence	54.8	47			49.1	63.9	
2016	Percentage of adults who have a regular health care provider	St. Lawrence	82.7	7 76.6	88.7	84.2	79	89.5	
	Percentage of adults who have taken a course or class to learn how to manage their						_		unreliable crude rate due to large standard error;
2016	chronic disease or condition	St. Lawrence	5.8	1.6	10.1	4.8	0	9.6	unreliable age-adjusted rate due to large standard error
	Percentage of adults who participated in leisure time physical activity in the past 30								
2016	days	St. Lawrence	70	63.6	76.3	69.2	62.4	76.1	
	Percentage of adults who report consuming less than one fruit and less than one								
2016	vegetable daily (no fruits or vegetables)	St. Lawrence	32.1	25.2	2 39	32.3	24.7	40	

			01.	CI 050/ I	CL 050/ 11		SI 050/ I A	SI 050/ II A	
Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate		CL 95% Upper, Age adjusted	Notes
						0			
									unreliable crude rate due to large standard error;
2016	Percentage of adults who use e-cigarettes	St. Lawrence	3.6	1.4	5.8	3.5	1.1	5.9	unreliable age-adjusted rate due to large standard error
2015	Percentage of adults with an annual household income less than \$25,000 who are	6	25.4						unreliable crude rate due to large standard error;
2016	obese	St. Lawrence	35.1	20.9	49.3	32.9	20.7	45	unreliable age-adjusted rate due to large standard error
	Developed find the other conditions that the character than \$25,000 days								and Palifer and a set of the form of the dead area.
2016	Percentage of adults with annual household income less than \$25,000 who are current smokers	Ch Lauranaa	23.5	10.8	26.3	24.7	13.7	25.0	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with arthritis	St. Lawrence St. Lawrence	32.6	26.7			25.5	35.6	
2010	Percentage of adults with artifitis  Percentage of adults with cardiovascular disease (heart attack, coronary heart	3t. Lawrence	32.0	20.7	36.3	30.9	23.3	30.3	
2016	disease, or stroke)	St. Lawrence	10.5	7.1	13.9	9.7	6.5	12.9	
2016	Percentage of adults with COPD	St. Lawrence	12.4				7.8		
2016	Percentage of adults with correct asthma	St. Lawrence	15				10.1	20.9	
	Percentage of adults with diagnosed high blood pressure taking high blood pressure								age-adjusted rate not defined due to empty age-
2016	medication	St. Lawrence	83.4	76.4	90.5	5			adjustment cells
2016	Percentage of adults with disability (based on 6 ACA disability questions)	St. Lawrence	26	20.5	31.5	24.1	18.9	29.2	
									unreliable crude rate due to large standard error;
2016	Percentage of adults with disability who are obese	St. Lawrence	40.3	29	51.7	31.2	18.6	43.8	unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with disability who smoke	St. Lawrence	17.8	9.1	26.5	24.5	9	39.9	unreliable age-adjusted rate due to large standard error
	Percentage of adults with disability with cardiovascular disease (heart attack,								
2016	coronary heart disease, or stroke)	St. Lawrence	26.8				8.2		unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with food security over the last 12 months	St. Lawrence	79.1	72.6	85.6		71.8		
2016	Percentage of adults with physician diagnosed diabetes	St. Lawrence	11.3				6.7		
2016	Percentage of adults with physician diagnosed high blood pressure.	St. Lawrence	35.6				27.3	38	
2016	Percentage of adults with physician diagnosed prediabetes	St. Lawrence	7.2	4.1	10.3	7.8	4.3	11.2	
2016	Percentage of adults with poor mental health for 14 or more days in the last month	St. Lawrence	9.8	5.9	13.7	10	5.9	14.1	
2015	Described for the soul in the state of the s	6							and the second state of th
2016	Percentage of cigarette smoking among adults who report poor mental health	St. Lawrence							rates suppressed due to small sample size
	Percentage of men aged 55-69 receiving shared decision making for PSA screening (discussed the advantages and disadvantages of the PSA test with a health								unreliable crude rate due to large standard error; age- adjusted rate not defined for indicators with an age
2016	professional)	St. Lawrence	29.4	17.3	41.5				restriction
2010	professional)	3t. Lawrence	25.4	17.3	41.3	1			restriction
2016	Percentage of women aged 18-44 years who had dental visit within the last year	St. Lawrence							rates suppressed due to small sample size
2010	Percentage of women aged 18-44 years whose health care provider discussed	St. Edwichte							races suppliessed due to small sample size
2016	planning a healthy pregnancy	St. Lawrence							rates suppressed due to small sample size
	, , , , , , , , , , , , , , , , , , ,								age-adjusted rate not defined for indicators with an age
2016	Percentage of women aged 18-64 years with health care coverage	St. Lawrence	99.5	98.7	100				restriction
	, ,								
	Percentage of women aged 21-29 years receiving cervical cancer screening based on								
	2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test								
2016	plus HPV test within past five years for women aged 30-65)	St. Lawrence							rates suppressed due to small sample size
	Percentage of women aged 21-65 years receiving cervical cancer screening based on							1	unreliable crude rate due to large standard error; age-
	2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test							1	adjusted rate not defined for indicators with an age
2016	plus HPV test within past five years for women aged 30-65)	St. Lawrence	87	75.9	98	3			restriction
	Percentage of women aged 50-74 years receiving breast cancer screening based on								age-adjusted rate not defined for indicators with an age
2016	recent guidelines	St. Lawrence	87.3	80.8	93.8	3			restriction
								1	
2016	Women aged 18-44 years who saw a doctor for a routine checkup within the last year	St. Lawrence						l	rates suppressed due to small sample size