

Behavioral Risk Factor Surveillance System (BRFSS) Health Indicators by County

Data from the 2016 Behavioral Risk Factor Surveillance System were used to generate percentages of non-institutionalized adult (18+) NYS residents for various health indicators for a range of geographies.

<https://health.data.ny.gov/Health/Behavioral-Risk-Factor-Surveillance-System-BRFSS-H/jsy7-eb4n>

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury	Warren	26.3	12	40.6				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults reporting a depressive disorder	Warren	16.7	12.3	21.2	18.3	12.9	23.8	
2016	Percentage of adult binge drinking during the past month	Warren	18.8	14.5	23.1	20.9	15.7	26.2	
2016	Percentage of adults aged 18+ years with flu immunization in the past year	Warren	37	31.8	42.2	32.9	27.8	37.9	
2016	Percentage of adults aged 18-24 years who are current smokers	Warren							rates suppressed due to small sample size
2016	Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year	Warren	67.8	61	74.6				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 18-64 years with healthcare coverage	Warren	93.4	90.3	96.6				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 50-64 years receiving colorectal cancer screening based on recent guidelines	Warren	68.8	59.7	78				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	Warren	75.1	68.6	81.7				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with at least one fall in the past 12 months	Warren	24.9	17.2	32.7				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with flu immunization in the past year	Warren	61	52.1	69.9				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with pneumococcal immunization	Warren	78.1	70.2	85.9				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults experiencing housing insecurity in the past 12 months	Warren	29.7	23.8	35.7	34.3	27.4	41.2	
2016	Percentage of adults overweight or obese	Warren	65.6	60	71.2	65.4	58.9	71.9	
2016	Percentage of adults reporting 14 or more days of poor physical health	Warren	11.4	8.1	14.6	10.9	7.4	14.4	
2016	Percentage of adults reporting driving under the influence in the past 30 days	Warren							rates suppressed due to small sample size
2016	Percentage of adults reporting poor self-reported health	Warren	2.6	0.8	4.4	2.9	0.4	5.3	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults who are current smokers	Warren	23.2	18	28.4	26.4	20.6	32.2	
2016	Percentage of adults who are obese	Warren	29.2	24.2	34.1	27.9	22.7	33.1	
2016	Percentage of adults who consume one or more sugary drinks daily	Warren	30	24.1	35.8	34.8	28.3	41.3	
2016	Percentage of adults who did not receive medical care because of cost	Warren	9.9	6.4	13.4	11	6.7	15.3	
2016	Percentage of adults who had a dentist visit within the past year	Warren	69.7	64.4	74.9	68.8	62.8	74.8	
2016	Percentage of adults who had a test for high blood sugar or diabetes within the past three years	Warren	56.9	51	62.8	52.6	46.2	59.1	
2016	Percentage of adults who have a regular health care provider	Warren	85.9	82	89.8	82.9	77.7	88.2	
2016	Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition	Warren	4.7	2.1	7.3	4	1.2	6.7	unreliable age-adjusted rate due to large standard error
2016	Percentage of adults who participated in leisure time physical activity in the past 30 days	Warren	79.6	75.1	84.1	79.8	74.5	85	
2016	Percentage of adults who report consuming less than one fruit and less than one vegetable daily (no fruits or vegetables)	Warren	27.6	22.1	33.2	29.4	23	35.8	
2016	Percentage of adults who use e-cigarettes	Warren	4.4	1.8	7	4.6	1.8	7.3	unreliable age-adjusted rate due to large standard error

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Percentage of adults with an annual household income less than \$25,000 who are obese	Warren	31.2	20	42.4	29.7	19.4	40.1	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with annual household income less than \$25,000 who are current smokers	Warren	33.2	20.4	46.1	34.8	25.2	44.4	unreliable crude rate due to large standard error
2016	Percentage of adults with arthritis	Warren	33.3	28.2	38.4	28.9	23.7	34.2	
2016	Percentage of adults with cardiovascular disease (heart attack, coronary heart disease, or stroke)	Warren	6	4	7.9	4.6	3	6.2	
2016	Percentage of adults with COPD	Warren	6	3.7	8.2	5.4	3.1	7.8	
2016	Percentage of adults with current asthma	Warren	10.4	7.1	13.8	11.6	7.2	16	
2016	Percentage of adults with diagnosed high blood pressure taking high blood pressure medication	Warren	72.5	63.2	81.7	44.2	36	52.3	small age-adjustment cell(s) (n = 2)
2016	Percentage of adults with disability (based on 6 ACA disability questions)	Warren	23	18.3	27.8	20.5	15.4	25.6	
2016	Percentage of adults with disability who are obese	Warren	43.5	31.7	55.3	42.5	28.1	56.8	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with disability who smoke	Warren	31.6	19.1	44	36	24.4	47.6	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with disability with cardiovascular disease (heart attack, coronary heart disease, or stroke)	Warren	12	5.7	18.2	12.2	2.8	21.6	unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with food security over the last 12 months	Warren	84.1	80	88.2	82.9	78.3	87.4	
2016	Percentage of adults with physician diagnosed diabetes	Warren	9.6	6.5	12.7	7.6	5.2	10.1	
2016	Percentage of adults with physician diagnosed high blood pressure.	Warren	36.5	30.9	42	31.9	26.2	37.6	
2016	Percentage of adults with physician diagnosed prediabetes	Warren	6.8	3.9	9.6	5.7	3.4	8	
2016	Percentage of adults with poor mental health for 14 or more days in the last month	Warren	10	6.4	13.7	12	7.4	16.7	
2016	Percentage of cigarette smoking among adults who report poor mental health	Warren							rates suppressed due to small sample size
2016	Percentage of men aged 55-69 receiving shared decision making for PSA screening (discussed the advantages and disadvantages of the PSA test with a health professional)	Warren	36.4	22.4	50.5				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-44 years who had dental visit within the last year	Warren	64.8	47.6	82.1				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy	Warren							rates suppressed due to small sample size
2016	Percentage of women aged 18-64 years with health care coverage	Warren	98.2	96.5	99.9				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 21-29 years receiving cervical cancer screening based on 2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test plus HPV test within past five years for women aged 30-65)	Warren							rates suppressed due to small sample size
2016	Percentage of women aged 21-65 years receiving cervical cancer screening based on 2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test plus HPV test within past five years for women aged 30-65)	Warren	87.7	80.4	95				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines	Warren	87.7	81.3	94.1				age-adjusted rate not defined for indicators with an age restriction
2016	Women aged 18-44 years who saw a doctor for a routine checkup within the last year	Warren	68.7	51.4	85.9				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction