

Behavioral Risk Factor Surveillance System (BRFSS) Health Indicators by County

Data from the 2016 Behavioral Risk Factor Surveillance System were used to generate percentages of non-institutionalized adult (18+) NYS residents for various health indicators for a range of geographies.

<https://health.data.ny.gov/Health/Behavioral-Risk-Factor-Surveillance-System-BRFSS-H/jsv7-eb4n>

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Among adults aged 65+ years with at least one fall in the past 12 months, percent that resulted in injury	Washington	29.6	16.8	42.4				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults reporting a depressive disorder	Washington	16.2	11.4	21	17.2	10.9	23.5	
2016	Percentage of adult binge drinking during the past month	Washington	19.1	13.8	24.5	21.7	15.5	27.8	
2016	Percentage of adults aged 18+ years with flu immunization in the past year	Washington	41.5	35.2	47.8	39.1	31.8	46.5	
2016	Percentage of adults aged 18-24 years who are current smokers	Washington							rates suppressed due to small sample size
2016	Percentage of adults aged 18-64 years who saw a doctor for a routine checkup within the last year	Washington	70.5	63.7	77.3				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 18-64 years with healthcare coverage	Washington	93.7	89.7	97.7				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 50-64 years receiving colorectal cancer screening based on recent guidelines	Washington	63.8	54	73.5				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 50-75 years receiving colorectal cancer screening based on recent guidelines	Washington	69	61.4	76.5				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with at least one fall in the past 12 months	Washington	31.3	23.4	39.2				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with flu immunization in the past year	Washington	55.8	46.8	64.9				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults aged 65+ years with pneumococcal immunization	Washington	75.6	67.6	83.6				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of adults experiencing housing insecurity in the past 12 months	Washington	32.2	25.8	38.6	35.9	28.1	43.7	
2016	Percentage of adults overweight or obese	Washington	71.3	65.7	76.8	68.5	61.6	75.5	
2016	Percentage of adults reporting 14 or more days of poor physical health	Washington	15.8	11.2	20.5	16.1	10.6	21.6	
2016	Percentage of adults reporting driving under the influence in the past 30 days	Washington							rates suppressed due to small sample size
2016	Percentage of adults reporting poor self-reported health	Washington	4.8	1.9	7.6	4.2	2	6.5	
2016	Percentage of adults who are current smokers	Washington	22.3	16.8	27.8	25.3	18.8	31.8	
2016	Percentage of adults who are obese	Washington	40.2	33.8	46.5	40.6	33.2	48.1	
2016	Percentage of adults who consume one or more sugary drinks daily	Washington	31.2	24.7	37.7	35.4	28.1	42.7	
2016	Percentage of adults who did not receive medical care because of cost	Washington	10.7	6.3	15	10.6	6.4	14.8	
2016	Percentage of adults who had a dentist visit within the past year	Washington	66.5	60.8	72.2	65.1	58.3	71.8	
2016	Percentage of adults who had a test for high blood sugar or diabetes within the past three years	Washington	55.5	48.7	62.2	55.4	49	61.9	
2016	Percentage of adults who have a regular health care provider	Washington	86.4	81.4	91.3	84.3	78.9	89.6	
2016	Percentage of adults who have taken a course or class to learn how to manage their chronic disease or condition	Washington	4.8	2.4	7.2	4.8	1.8	7.9	unreliable age-adjusted rate due to large standard error
2016	Percentage of adults who participated in leisure time physical activity in the past 30 days	Washington	75.2	70.3	80.2	77.5	72.6	82.5	
2016	Percentage of adults who report consuming less than one fruit and less than one vegetable daily (no fruits or vegetables)	Washington	27.4	21.6	33.2	26.9	20.3	33.5	
2016	Percentage of adults who use e-cigarettes	Washington	4.1	0.9	7.4	4.6	1.3	7.9	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Percentage of adults with an annual household income less than \$25,000 who are obese	Washington	45.4	32.1	58.8	52.4	41.3	63.6	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with annual household income less than \$25,000 who are current smokers	Washington	27.8	16.8	38.9	28.1	16.9	39.2	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with arthritis	Washington	33.8	28.5	39.2	28.7	23.9	33.5	
2016	Percentage of adults with cardiovascular disease (heart attack, coronary heart disease, or stroke)	Washington	10.6	7	14.2	8.2	5.6	10.7	
2016	Percentage of adults with COPD	Washington	7	4.5	9.6	5.9	3.7	8.2	
2016	Percentage of adults with current asthma	Washington	9.3	5.9	12.8	8.2	5.1	11.2	
2016	Percentage of adults with diagnosed high blood pressure taking high blood pressure medication	Washington	72	61.1	82.8	51.4	41.3	61.4	unreliable crude rate due to large standard error; unreliable age-adjusted rate due to large standard error
2016	Percentage of adults with disability (based on 6 ACA disability questions)	Washington	24	19.1	29	21.8	15.9	27.6	
2016	Percentage of adults with disability who are obese	Washington	55.4	44.3	66.5	57.8			unreliable crude rate due to large standard error; variability of age-adjusted rate not defined due to age-adjustment cells with 1 observation
2016	Percentage of adults with disability who smoke	Washington	33	20.1	45.9	38.2			unreliable crude rate due to large standard error; variability of age-adjusted rate not defined due to age-adjustment cells with 1 observation
2016	Percentage of adults with disability with cardiovascular disease (heart attack, coronary heart disease, or stroke)	Washington	21.8	11.4	32.2	15.5			unreliable crude rate due to large standard error; variability of age-adjusted rate not defined due to age-adjustment cells with 1 observation
2016	Percentage of adults with food security over the last 12 months	Washington	76.6	70.7	82.6	72.7	65.2	80.1	
2016	Percentage of adults with physician diagnosed diabetes	Washington	13.5	10	17	11.2	8.5	14	
2016	Percentage of adults with physician diagnosed high blood pressure.	Washington	35.7	29.6	41.8	34.5	27.9	41.1	
2016	Percentage of adults with physician diagnosed prediabetes	Washington	6.6	3.6	9.5	6.5	3.5	9.5	
2016	Percentage of adults with poor mental health for 14 or more days in the last month	Washington	12.2	8.2	16.2	13.1	8.8	17.4	
2016	Percentage of cigarette smoking among adults who report poor mental health	Washington	51	33.1	68.8	41.8			unreliable crude rate due to large standard error; variability of age-adjusted rate not defined due to age-adjustment cells with 1 observation
2016	Percentage of men aged 55-69 receiving shared decision making for PSA screening (discussed the advantages and disadvantages of the PSA test with a health professional)	Washington	37.2	23.9	50.4				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-44 years who had dental visit within the last year	Washington	70.1	54.8	85.3				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-44 years whose health care provider discussed planning a healthy pregnancy	Washington	46	27.9	64				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 18-64 years with health care coverage	Washington	94.5	90.2	98.9				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 21-29 years receiving cervical cancer screening based on 2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test plus HPV test within past five years for women aged 30-65)	Washington							rates suppressed due to small sample size
2016	Percentage of women aged 21-65 years receiving cervical cancer screening based on 2012 guidelines (Pap test within past three years for women aged 21-65, or Pap test plus HPV test within past five years for women aged 30-65)	Washington	87.8	79.8	95.9				age-adjusted rate not defined for indicators with an age restriction
2016	Percentage of women aged 50-74 years receiving breast cancer screening based on recent guidelines	Washington	78.6	69.9	87.4				age-adjusted rate not defined for indicators with an age restriction

Year	Health Indicator	County	Crude Rate	CL 95% Lower, Crude	CL 95% Upper, Crude	Age-adjusted Rate	CL 95% Lower, Age-adjusted	CL 95% Upper, Age-adjusted	Notes
2016	Women aged 18-44 years who saw a doctor for a routine checkup within the last year	Washington	75.4	62.8	88.1				unreliable crude rate due to large standard error; age-adjusted rate not defined for indicators with an age restriction